



AD	-A19	94 717 <u>:</u>	TEPPIT DOCU	MENTATION	PAGE U	III. C	ILE CUPY
, , , ,			ECTE	16. RESTRICTIVE	MARKINGS		
28. SECURITY CLASSIFICATION AUTICIPITY MAY 1 7 1988				3. DISTRIBUTION / AVAILABILITY OF REPORT			
2b. DECLASSIFICATION / DOWNGRACING SCHEDULE				Approved for public release; distribution unlimited.			
4. PERFORMING ORGANIZATION REPORT NUMBER				5. MONITORING ORGANIZATION REPORT NUMBER(S)  ARO 22109.6-CH			
6c. ADDRESS (City, State, and ZIP Code)				7b. ADDRESS (City, State, and ZIP Code)			
Tucson, AZ 85721				P. O. Box 12211 Research Triangle Park, NC 27709-2211			
ORGANIZA	· - ·	onsoring rch Office	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DAAG29-85-K-0068			
	(City, State, and			10. SOURCE OF FUNDING NUMBERS			
P. O. Box 12211 Research Triangle Park, NC 27709-2211				PROGRAM ELEMENT NO.	PROJECT NO.	TASK NO.	WORK UNIT ACCESSION NO.
Polymer		f Azaethylenes	s (Imines) and A	za-1,3-Diene	es. Potenti	al React	ive Monomers
13a. TYPE OF		13h TIME CO	OVERED 1/85 TO 2/29/8	14. DATE OF REPORT (Year Month, Day) 15. PAGE SOUNT			
	entary NOTA	The view,	opinions and/or t be construed a	findings co			
17	COSATI	CODES	18. SUBJECT TERMS (	nomers, Polymerization, Butadienes,  1 Reactive Monomers			
FIELD	GROUP	SUB-GROUP					
19 ABSTRACT	T (Continue co	reverse if necessari	and identify by block i				
	Severa	al types of ne	ew imine monomer	s have been			
p	olymerizal	bility has the	en been investig	ated in diff	erent condi	cions.	ine main
С	lasses of	imines studie	ed include 1-aza	-1,3-butadie	nes with an	electro	n
d	onating o	r electron-acc	cepting substitu	ent on nitro	gen, 2-aza-	1,3-buta	dienes
w	ith a sub	stituent in th	ne l-position, i	mines with t	hree electr	on-accep	ting.
S	ubstituen	ts, and imines	s with only one	electron-acc	epting subs	stituent.	
	ITION/AVAILAE SSIFIED/UNLIMI	BILITY OF ABSTRACT	RPT. DTIC USERS		CURITY CLASSIFIC	CATION	
22a NAME C	OF RESPONSIBL	E INDIVIDUAL			(Include Area Cod	le) 22c. OFF	CE SYMBOL
DD FORM 1	473, 84 MAR	83 AI	PR edition may be used us	ntil exhausted	SECURITY	CLASSIFICAT	TION OF THIS PAGE

Polymerization of Azaethylenes (Imines) and Aza-1,3-dienes. Potential Reactive Monomers.

FINAL REPORT

H. K. Hall, Jr.

April 1, 1988

U.S. Army Research Office

Grant DAAG29-85-K-0068

University of Arizona

Approved for Public Release; distribution unlimited

## Statement of Problem

Although a great deal of chemistry is known about the polymerization of C=C and C=0 compounds, very little is known about the polymerization of C=Ncompounds. The literature does contain tantalizing hints of the ability of azaethylenes (imines) to polymerize. Thus, N-alkylazaethylenes are known to cyclotrimerize. However, before our work, no systematic study has been carried out.

We shall call imines "azaethylenes" to point out their relationship to vinyl monomers. The aza-1,3-dienes are then equivalents of butadienes.

> Unamiounced Justification Distribution/ Availability Codes Avail and/or Dist Special

Accession For NTIS GRA&I DTIC TAB

### Results

Several types of new imine monomers have been synthesized and their polymerizability has then been investigated in different conditions. The main classes of imines studied include 1-aza-1,3-butadienes with an electron donating or electron-accepting substituent on nitrogen, 2-aza-1,3-butadienes with a substituent in the 1-position, imines with three electron-accepting. Substituents, and imines with only one electron-accepting substituent.

### 1-Azabutadienes

We began our work with a study of the synthesis and polymerization of 1-azabutadienes:

We synthesized various 1-substituted 1-azabutadienes via pyrolysis of cyclopentadiene adducts. We found that the substituent on N and the type of initiator played major roles in bringing about 4,1 polymerization of 1-azabutadienes. Donor groups on N such as CH<sub>3</sub>O-, (CH<sub>3</sub>)<sub>2</sub>N-; C<sub>6</sub>H<sub>5</sub>-with electrophilic initiators did not give cationic polymerization, but only stable ammonium salts. But acceptor groups (C<sub>6</sub>H<sub>5</sub>-,-COOC<sub>2</sub>H<sub>5</sub>) with anionic initiators gave high yields of high polymer with 4,1 structure (2).

$$+CH_{2}-CH=CH-N+_{M}$$

$$+CH_{2}-CH+_{M}$$

$$CH=N-_{R}$$

$$4.1- structure$$

$$4.3- structure$$

$$R=C_{6}H_{5}^{-},-COOC_{2}H_{5}$$

With <u>free radical</u> initiators, 1-azabutadienes with donor or acceptor groups on nitrogen gave indifferent results. One problem we encountered was

that N-carboethoxy-1-azabutadiene tends to dimerize in these conditions  $(70^{\circ})$ . This unwanted hetero-Diels-Alder reaction of the C=N-C=O unit effectively competes with the radical polymerization. (The anionic polymerization mentioned above proceeds faster at lower temperature.) We had earlier demonstrated this ability of a -COOR acceptor group in conjugation with C=N group to undergo hetero-Diels-Alder for triethyl ethyleneiminetricarboxylate.

### 2-Azabutadienes

Several 1-substituted-2-azabutadienes have also been synthesized via pyrolysis. The substituents were phenyl, 2,4,6-trimethylphenyl and t-butyl. Only the first one led to low molecular weight polymer. Thus, we showed that polymerization involving propagation in the sense  $\longrightarrow$  \* + N=C  $\longrightarrow$  N-C\*, where \* = +, ., or - is not suitable.

$$R = -C_6H_5$$
,  $-C_6H_2Me_3$ ,  $-CMe_3$ 

## Azaethylenes (Imines)

We have devised, based on literature suggestions, a new synthesis of azaethylenes with three acceptor groups.

So far, we have studied:

$$\begin{bmatrix} NC - N = C \\ CN \end{bmatrix}$$

$$\begin{bmatrix} NC - N = C \\ COOCH_3 \end{bmatrix}$$

The  $C_4N_4$  is very reactive. Attempted isolation gives only oligomeric material, even when acidic stabilizers or freeze-drying are employed. The cyanoester monomer is more stable in solution and its spectra can be obtained. Thus, ester groups instead of cyano groups on carbon offer somewhat lower reactivity and hope of isolation and purification. Nevertheless, again isolation does not succeed. Conclusive evidence for the presence of both of these highly reactive azaethylene monomers was obtained by trapping them as Diels-Alder cycloadducts with cyclopentadiene:

Upon mixing with p-methoxystyrene, these two highly electrophilic azaethylenes spontaneously formed strictly alternating copolymers.

$$\begin{bmatrix} NC - N = C & N \\ N & - C & - CH_2 - CH \\ N & CN & CN \\ N & CN & CN & COH. \end{bmatrix}$$

Turning to less reactive imines, we turned to azaethylenes with only an electron-accepting substituent on N. In order to avoid tautomerization to enimides, the substituents on carbon should not have any  $\beta$ -hydrogens.

The following imines were selected:

The N-cyano and N-sulfonyl imines were synthesized using silylated reagents. For example.

The N-carboethoxy imines were synthesized by reaction of the aldehyde with ethyl carbonate:

$$\emptyset$$
 CH=O + H<sub>2</sub>N - COOEt  $\longrightarrow$   $\emptyset$ -CH = N - COOEt

The N-cyanoimines both polymerized in high yield under anionic or free radical conditions, but molecular weights were low. The other three imines did not polymerize. The low steric requirement of =N-CN (linear) versus = N-COOEt or = N - SO<sub>2</sub>Me appears responsible.

#### Conclusions

To sum up our USARO work, we have made great progress in understanding how to design and polymerize azaethylene monomers:

- We have designed new general synthesis of N-acceptor-azaethylene monomers.
- 2. Suitable acceptor groups on N include CN and COOR.
- 3. Two acceptor groups on carbon have led to too reactive monomers, although they can be detected in solution, and monomers with less powerful acceptor groups may still be isolable.
- Donor substituents on carbon should be aryl or t-butyl. Alkyls with  $\beta$ -H are unsuitable because of tautomerization. Two hydrogens will lead to too reactive monomers.
- 5. Anionic and free radical polymerization are the methods of choice.
- 6. The C=N-Acc monomers and polymerization resemble the anionic polymerization of aldehydes.

The polymers obtained so far have been white solids with tg 50-125°, in general, soluble in organic solvents, and suitable for binder materials.

## Publications under USARO Sponsorship

- 1. T. Kitayama and H.K. Hall, Jr., "Synthesis and Polymerization of 1-Azabutadiene Monomers," <u>Macromolecules</u>, <u>20</u>, 1451 (1987).
- 2. J. Oku, A.B. Padias, H.K. Hall, Jr. and A.J. East, "Poly (3-Buty1-3-ethyl Malonimide)," Macromolecules, 20, 2314 (1987).
- 3. T. Kitayama, A.B. Padias, and H.K. Hall, Jr., "On the Polymerization of 1-substituted 1-Aza-1,3-butadienes," Polym. Bull., 17, 417 (1987).
- 4. M. Ramezanian, F.D. Saeva and H.K. Hall, Jr., "A New Azacyanocarbon,  $C_4N_4$ ; Tricyanomethanimine," <u>Tetrah. Lett</u>. (in press).
- 5. J.B. Kim and H.K. Hall, Jr. "Synthesis and Polymerization of 1-Carboethoxy-3-methyl-1-aza-1,3-butadiene," <u>Macromolecules</u> (in press).

- J.B. Kim, A.B. Padias and H.K. Hall, Jr., "Synthesis and Polymerization of Electron-Deficient Imines," (in preparation).
- M. Ramezanian and H.K. Hall, Jr., "Synthesis and Polymerization of N-Cyanoimines", (in preparation).

# Scientific Personnel

H.K. Hall, Jr. Principal Investigator, 3 summer months

T. Kitayama Postdoctoral, 1 year

J. B. Kim Postdoctoral, 1 year

J. I.Oku Postdoctoral, 6 months

J. I. Weinschenk Graduate Student, 6 months

M. Ramezanian Graduate Student, 18 months